

SCHWINGER EFFECT IN STRING THEORY ^a

C.BACHAS

*Centre de Physique Théorique, Ecole Polytechnique
91128 Palaiseau, France*

ABSTRACT

I discuss the quantum instability of an electric field in a theory of open strings.

One of the few available hints on the interplay between quantum theory and gravity is the Hawking emission of a black hole ¹. This can be described as pair creation in the background gravitational field. Of the two produced particles one stays trapped inside the horizon, while the other escapes to infinity where it emerges with a thermal spectrum at an effective temperature $T_H = (8\pi GM)^{-1}$. In deriving his famous result Hawking assumed a Schwarzschild form for the metric, and minimally-coupled non-self-interacting quantum fields. In string theory we expect two kinds of corrections to this result: (a) Quantum or string-loop effects, which are for sure important for small black holes of mass $M \simeq M_{Planck} \simeq 1/\sqrt{\alpha'} g_{string}$, and (b) σ -model or α' corrections, which when $g_{string} \ll 1$ can become important much earlier, for $M \simeq M_{Planck}/g_{string} \simeq 1/\sqrt{\alpha'} g_{string}^2$. Assuming the spectrum stays thermal, these latter corrections will presumably modify the expression for the temperature ² by replacing $GM \rightarrow GM \times f(GM/\sqrt{\alpha'})$, where $f(y) \simeq 1 + o(\frac{1}{y^2})$ for large y . Knowledge of the function f , which parametrizes the effects of the non-minimal coupling of strings to gravity, could be a crucial first step before addressing the paradoxes related to black-hole evaporation.

A similar but simpler problem arises when one considers the quantum instability of strong electromagnetic fields. Schwinger's field-theoretic calculation ³, for minimally-coupled non-self-interacting particles, gives the following result for the rate of pair production per unit volume and time:

$$w = \frac{2J+1}{8\pi^3} \sum_{k=1}^{\infty} (-)^{(2J+1)(k+1)} \left(\frac{eE}{k}\right)^2 \exp\left(-\frac{k\pi m^2}{|eE|}\right). \quad (1)$$

Here E is the constant external electric field, e and m are the charge and mass of the produced particles, and $J = 0$ or $\frac{1}{2}$ is their spin. This rate, obtained as the imaginary part of the one-loop vacuum-energy density, vanishes non-perturbatively

^aProceedings of the Rome Workshop on *Quantum Gravity, String Theory and Grand Unification*, Rome (September 1992).

fast with the external field, showing that pair-production is a quantum-tunneling phenomenon. The potential barrier has, roughly-speaking, a height of order $2m$ and a width of order m/eE .

This calculation can be also done exactly, if point-particles are replaced by open strings ⁴. The key observation ⁵ is that the effect of a constant electric field $F^{01} = E$ is equivalent to an orbifold-like twist of the light-like coordinates $X^\pm = \frac{1}{\sqrt{2}}(X^0 \pm X^1)$, by an imaginary angle $\pm i\epsilon$, where

$$\epsilon = \frac{1}{\pi} [\text{arcth}(\pi e_1 E) + \text{arcth}(\pi e_2 E)] . \quad (2)$$

Here e_1 and e_2 are the two charges at the string endpoints, and we have set $2\alpha' = 1$. Taking into account the above twist, and the anomalous commutation relation of the center-of-mass coordinates $[x^+, x^-] = \frac{-i}{(e_1 + e_2)E}$, one arrives ⁴ at the following expression for the annulus contribution to the vacuum energy density of oriented bosonic strings :

$$\mathcal{F} = \int_0^\infty \frac{dt}{t} (2\pi^2 t)^{-\frac{D}{2}} Z(t) f_A(t, e_1 E, e_2 E) , \quad (3)$$

where $Z(t) = \sum_{states} e^{-\frac{\pi t}{2} m_S^2}$ is the usual partition function of an open string, D is the number of non-compact space-time dimensions, and the entire field dependence is contained in the correction factor

$$f_A = \frac{\pi(e_1 + e_2)E t e^{-\pi t \epsilon^2/2}}{2 \sin(\pi t \epsilon/2)} \prod_{n=1}^{\infty} \frac{(1 - e^{-\pi t n})^2}{(1 - e^{-\pi t(n+i\epsilon)})(1 - e^{-\pi t(n-i\epsilon)})} . \quad (4)$$

This factor goes to one in the zero-field limit, and has evenly-spaced simple poles at $t = \frac{2k}{|\epsilon|}$ ($k = 1, 2, \dots$) on the integration axis. These poles give in turn rise to an imaginary part of $\mathcal{F}$, and hence to a string pair-production rate

$$w_{bos} = \frac{1}{2(2\pi)^{D-1}} \sum_{states \ S} \frac{(e_1 + e_2)E}{\epsilon} \sum_{k=1}^{\infty} (-)^{k+1} \left(\frac{|\epsilon|}{k}\right)^{D/2} \exp\left(-\frac{\pi k}{|\epsilon|} (m_S^2 + \epsilon^2)\right) . \quad (5)$$

The above result stays valid for unoriented strings ⁴, for which the projection onto physical states is achieved by adding to the annulus the contribution of the Möbius-strip ⁶. For the fermionic string, on the other hand, a straightforward extension of the calculation ⁴ gives

$$w_{sustr} = \frac{1}{2(2\pi)^{D-1}} \sum_{states \ S} \frac{(e_1 + e_2)E}{\epsilon} \sum_{k=1}^{\infty} (-)^{(k+1)(a_S+1)} \left(\frac{|\epsilon|}{k}\right)^{D/2} \exp\left(-\frac{\pi k m_S^2}{|\epsilon|}\right), \quad (6)$$

where $a_S = 0$ or 1 according to whether the state S belongs to the Neveu-Schwarz or the Ramond sector.

The string-pair-production rate, eqs.(5,6), reduces to Schwinger's result, eq.(1), in the weak-field limit, as has been noticed before by Burgess ⁷. The non-minimal electromagnetic coupling of strings manifests, however, itself for stronger E , and is essentially summarized by the replacement:

$$(e_1 + e_2)E \rightarrow \epsilon \simeq (e_1 + e_2)E \times [1 + o((E\alpha')^2)] . \quad (7)$$

The non-linear function ϵ , as well as the string-pair-production rate, diverge at the limiting field value $E_{lim} = 1/2\pi\alpha' \max(e_i)$. The existence of a limiting field has been noticed also at the classical level ^{8,5,7}, but the above result clearly establishes the responsible dissipation mechanism. Analogous results for strong gravitational, or non-abelian fields would be extremely interesting. The latter could in particular drastically modify our current semi-phenomenological ⁹ understanding of fragmentation in Q.C.D.

Before concluding let me note that while the string-structure seems to enhance the effects of an electric field, it moderates those of a magnetic field. A magnetic field, $F^{12} = B$, twists the complex space-coordinate $X = \frac{1}{\sqrt{2}}(X^1 + iX^2)$ by a real phase ⁵

$$\tilde{\epsilon} = \frac{1}{\pi} [\arctan(\pi e_1 B) + \arctan(\pi e_2 B)] , \quad (8)$$

which determines the spacing between Landau levels. For sufficiently massive string states this spacing reads $\tilde{\epsilon}/2m$. It reduces to the usual cyclotron frequency for weak B , but levels off towards a limiting value $1/4\alpha'm$ as the field becomes stronger. This behaviour can be understood as due to a maximum allowed centripetal acceleration. Notice that the degeneracy of any given Landau level is the same as for a point particle, since the commutation relation of the center-of-mass coordinates is not modified ⁵. Finally let me point out that the annulus amplitude in the presence of a B -field has no poles on the integration axis. Thus, as in point-particle field theory, a magnetic field cannot dissipate by spin-polarizing the vacuum.

This research was supported in part by EEC contracts SC1-0394-C and SC1-CT92-0792.

References

1. S.W. Hawking, *Nature* **248** (1974) 30 ; *Comm.Math.Phys.* **43** (1975) 199.
2. C.G.Callan, R.C.Myers and M.J.Perry, *Nucl. Phys.* **B311** (1988/89) 673.
3. J.Schwinger, *Phys.Rev.* **82** (1951) 664.
4. C.Bachas and M.Porrati, *Phys.Lett.* **296B** (1992) 77.
5. A.Abouelsaood, C.G.Callan, C.R.Nappi and S.A.Yost, *Nucl. Phys.* **B280** [FS18] (1987) 599.
6. See for instance A.Sagnotti, in *Non-Perturbative Quantum Field Theory*, G.'t Hooft et al eds., (Plenum, 1988), p. 521.

7. C.P.Burgess, Nucl.Phys. **B294** (1987) 427.
8. E.S.Fradkin and A.A.Tseytlin, Phys.Lett. **163B** (1985) 123; V.V.Nesterenko, Int.J.Mod.Phys. **A4** (1989) 2627.
9. B.Andersson, G.Gustafson, G.Ingelman and T.Sjöstrand, Phys.Rep. **97** (1983) 33.